

Frank's constant in the hexatic phase

P. Keim, G. Maret*, and H.H. von Grünberg

Universität Graz, 8010 Graz, Austria

**Universität Konstanz, 78457 Konstanz, Germany*

(Dated: February 6, 2008)

Using video-microscopy data of a two-dimensional colloidal system the bond-order correlation function G_6 is calculated and used to determine the temperature-dependence of both the orientational correlation length ξ_6 in the isotropic liquid phase and the Frank constant F_A in the hexatic phase. F_A takes the value $72/\pi$ at the hexatic $\leftrightarrow$ isotropic liquid phase transition and diverges at the hexatic $\leftrightarrow$ crystal transition as predicted by the KTHNY-theory. This is a quantitative test of the mechanism of breaking the orientational symmetry by disclination unbinding.

PACS numbers: 64.70.Dv, 68.35.Rh, 82.70.Dd

The theory of melting in two dimensions (2d) developed by Kosterlitz, Thouless, Halperin, Nelson and Young (KTHNY-theory) suggests a two-stage melting from the crystalline phase to the isotropic liquid. The first transition at temperature T_m is driven by the dissociation of thermally activated dislocation pairs into isolated dislocations breaking the translational symmetry [1, 2]. The fluid phase directly above T_m still exhibits orientational symmetry and is called the hexatic phase. It may be viewed as an anisotropic fluid with a six-fold director [3, 4] which is characterized by a finite value of Frank's constant F_A , the elastic modulus quantifying the orientational stiffness. At the second transition at $T_i > T_m$, the dissociation of some of the dislocations into free disclinations destroys the orientational symmetry. Now, the fluid shows ordinary short-range rotational and positional order as it is characteristic of an isotropic liquid.

Following an argument given in [1, 4], T_m and T_i can be estimated using the defect interaction Hamiltonian H_d between a pair of disclinations ($d = disc$) and a pair of dislocations ($d = disl$) which for both defect pairs and at large distances goes like $H_d \sim c_d \log r$ with the dimensionless strength parameter c_d depending on the defect type. Defect dissociation is completed at a temperature where the thermally averaged pair distance $\langle r_d^2 \rangle$ diverges. Evaluating this expression for H_d one generally finds divergence if $c_d = 4$. The unbinding condition $c_d = 4$ translates into $\lim_{T \rightarrow T_m^-} \beta \mathcal{K}(T) a_0^2 = 16\pi$ for dislocation pairs ($\beta = 1/k_B T$, a_0 is lattice spacing) and into $\lim_{T \rightarrow T_i^-} \beta F_A(T) = 72/\pi$ for disclination pairs, where $\mathcal{K}$ is the Young's modulus of the crystal. Connecting thus the defect pair unbinding condition to the two transition temperatures T_i and T_m , two expressions are obtained that summarize the microscopic explanation of the KTHNY theory for two-stage melting.

In this Letter we study the temperature-dependence of Frank's constant of a 2D system in the hexatic phase. We first determine the hexatic $\rightarrow$ isotropic fluid transition temperature T_i and then check if Frank's constant takes the value $72/\pi$ at T_i , thus testing the KTHNY theory

and its prediction that disclination unbinding occurs at T_i . In addition, we analyze the divergence behavior of the orientational correlation length at T_i and of Frank's constant at T_m .

Different theoretical approaches invoking grain boundary induced melting [5, 6] or condensation of geometrical defects [7, 8] suggest one first order transition. However some simulations for Lennard-Jones systems indicate the hexatic phase to be metastable [9, 10]. The transition in hard-core systems seem to be first-order [12] probably due to finite-size effects [11]. Simulations with long-range dipole-dipole interaction clearly show second order behavior [13]. Experimental evidence for the hexatic phase has been demonstrated for colloidal systems [14, 15, 16, 17, 18, 19], in block copolymer films [20, 21], as well as for magnetic bubble arrays and macroscopic granular or atomic systems [22, 23, 24, 25, 26]. Still the order of the transitions is seen to be inconsistent. The observation of a phase equilibrium isotropic/hexatic [17, 21] and hexatic/crystalline [17] indicates two first order transitions. In our system we find two continuous transitions.

The experimental setup is essentially the same as in [27]. Spherical and super-paramagnetic colloids (diameter $d = 4.5 \mu m$) are confined by gravity to a water/air interface formed by a water drop suspended by surface tension in a top sealed cylindrical hole of a glass plate. The field of view has a size of $835 \times 620 \mu m^2$ containing typically up to $3 \cdot 10^3$ particles (out of $3 \cdot 10^5$ of the whole sample). A magnetic field $\vec{H}$ is applied perpendicular to the air/water interface inducing in each particle a magnetic moment $\vec{M} = \chi \vec{H}$. This leads to a repulsive dipole-dipole pair-interaction with the dimensionless interaction strength given by $\Gamma = \beta(\mu_0/4\pi)(\chi H)^2(\pi\rho)^{3/2}$. Here χ is the susceptibility per colloid while ρ is the 2d particle density and the average particle distance is $a = 1/\sqrt{\rho}$. The interaction strength can be externally controlled by means of the magnetic field H ; it can be interpreted as an inverse temperature and is the only parameter controlling the phase behavior of the system. For each Γ the coordinates of the colloids are recorded




FIG. 1: Structure factor $S(\vec{q})$ of our colloidal system at three different inverse temperatures Γ corresponding to the isotropic liquid ($\Gamma = 52.4$), the hexatic phase ($\Gamma = 59.6$) and the crystalline ($\Gamma = 61.0$) phase. The central cross is an artifact of the Fourier-transformation.

via video-microscopy (resolution of particle position $dr = 100 \text{ nm}$) and digital image processing over a period of $1 - 2 \text{ h}$ using a frame rate of 250 ms .

To set the stage we first visualize in Fig. (1) the three phases and their symmetries by plotting the structure factor

$$S(\vec{q}) = \frac{1}{N} \left\langle \sum_{\alpha, \alpha'} e^{-i\vec{q}(\vec{r}_\alpha - \vec{r}_{\alpha'})} \right\rangle, \quad (1)$$

as calculated from the positional data of the colloids for three different temperatures. Here, α, α' runs over all N particles in the field of view while $\langle \rangle$ denote the time average over 700 configurations. In the liquid phase, concentric rings appear having radii that can be connected to typical inter-particle distances. The hexatic phase, on the other hand, is characterized by six segments of a ring which arise due to the quasi long-range orientational order of the six-fold director [28]. In the crystalline phase the Bragg peaks of a hexagonal crystal show up with a finite width that is due to the quasi long-range character of the translational order.

To quantify the six-fold orientational symmetry the bond-order correlation function

$$G_6(r) = \langle \psi(\vec{r}) \psi^*(\vec{0}) \rangle, \quad (2)$$

is calculated with $\psi(\vec{r}) = \frac{1}{N_j} \sum_j e^{6i\theta_{ij}(\vec{r})}$. Here the sum runs over the N_j next neighbors of the particle i at position $\vec{r}$ and $\theta_{ij}(\vec{r})$ is the angle between a fixed reference axis and the bond of the particle i and its neighbor j . $\langle \rangle$ here denotes not only the ensemble average which is taken over all $N(N-1)/2$ particle-pair distances for each configuration (resolution $dr = 100 \text{ nm}$) but also the time average over 70 statistically independent configurations.

KTHNY theory predicts that

$$\begin{aligned} \lim_{r \rightarrow \infty} G_6(r) &\neq 0 && \text{crystal: long range order} \\ G_6(r) &\sim r^{-\eta_6} && \text{hexatic: quasi long range} \\ G_6(r) &\sim e^{-r/\xi_6} && \text{isotropic: short range} \end{aligned}$$

$\eta_6 < 1/4$ and takes the value $1/4$ right at $T = T_i$. All three regimes can be easily distinguished in Fig. (2) showing $G_6(r)$ for a few representative temperatures. Note, that $G_6(0)$ is not normalized to 1.

We next fit $G_6(r)$ to $r^{-\eta_6}$ and e^{-r/ξ_6} to extract η_6 and ξ_6 . The fits are performed for radii $r/a \in \{0..20\}$ [29]. To check for the characteristics of the orientational correlation function, the ratio of the reduced chi-square




FIG. 2: Orientational correlation function $G_6(r)$ as function of the inverse temperature Γ in a log-log plot. From top to bottom: Three curves for the crystalline phase showing the long-range orientational order ($\lim_{r \rightarrow \infty} G_6(r) \neq 0$), two curves showing the quasi long-range order of the hexatic phase ($G_6(r) \sim r^{-\eta_6(\Gamma)}$) and three curves showing the short-range order typical of the isotropic liquid ($G_6(r) \sim e^{-r/\xi_6(\Gamma)}$).




FIG. 3: Quantitative test for the long distance behavior of $G_6(r)$. For $\chi^2_{alg}/\chi^2_{exp} < 1$ the algebraic decay fits better.

χ^2 goodness-of-fit statistic of the algebraic (χ^2_{alg}) and exponential (χ^2_{exp}) fit is shown in Fig. (3) as a function of Γ for three different measurements. For melting, a crystal free of dislocations was grown at high Γ and then Γ was reduced in small steps. For each temperature step the system was equilibrated $1/2 h$ before data acquisition started. This was done at different densities: *melt_1* with average particle distance of $a = 11.8 \mu m$ and *melt_2* with $a = 14.8 \mu m$ containing 3200 respectively 2000 particles in the field of view. The measurement denoted *freeze* in Fig. (3) ($a = 11.8 \mu m$) started in the isotropic liquid phase and Γ was increased with an equilibration time of $1 h$ between the steps. For $\chi^2_{alg}/\chi^2_{exp} > 1$ an exponential decay fits better than the algebraic and vice versa for $\chi^2_{alg}/\chi^2_{exp} < 1$. We observe in Fig. (3) that the change in the characteristic appears at $\Gamma_i = 57.5 \pm 0.5$. This value is the temperature of the hexatic $\leftrightarrow$ isotropic liquid transition.

In the vicinity of the phase transition, approaching Γ_i from the isotropic liquid the orientational correlation length ξ_6 should diverge as [4],

$$\xi_6(\Gamma) \sim \exp\left(\frac{b}{|1/\Gamma - 1/\Gamma_i|^\nu}\right), \quad (3)$$

with b a constant and $\nu = 1/2$. This behavior is observed in Fig. (4a). ξ_6 indeed increases dramatically near $\Gamma_i = 57.5 \pm 0.5$ irrespective of whether the system is heated or cooled. Before discussing this feature we first address the finite size effect. To this end, we have computed $G_6(r)$ and ξ_6 for subsystems of different size ranging from $720 \times 515 \mu m^2$, $615 \times 405 \mu m^2$, $505 \times 300 \mu m^2$, $400 \times 190 \mu m^2$ to $390 \times 80 \mu m^2$. The resulting data-points are plotted as triangles in Fig. (4) and belong to the black filled squares which they converge to. No finite size effect is found for $\Gamma < 56$, but a considerable one at $\Gamma = 56.9$ close to Γ_i where we obviously need the full field of view to capture

the characteristic of the divergence. At $\Gamma = 58.0$ there is a huge finite size effect indicating that ξ_6 is much larger than the field of view. However, inside the hexatic phase, ξ_6 is no longer well defined as the decay is algebraic. We fit our data to eq. (3) in the range $49 < \Gamma < 57.5$ and find the critical exponent $\nu = 0.5 \pm 0.03$ and $\Gamma_i = 58.9 \pm 1.1$, a value which due to the finite-size effect is slightly larger than Γ_i obtained from Fig. (3).

The exponent η_6 is related to Frank's constant F_A [4]

$$\eta_6(\Gamma) = \frac{18k_B T}{\pi F_A(\Gamma)}. \quad (4)$$

So the critical exponent $\eta_6(\Gamma_i) = 1/4$ corresponds to $\beta F_A(\Gamma_i) = 72/\pi$ at the hexatic $\leftrightarrow$ liquid transition. This quantity is plotted in Fig. (4b). Indeed, F_A crosses the value $72/\pi$ at $\Gamma_i = 57.5 \pm 0.5$ exactly at that temperature which in Fig. (3) has been independently determined to




FIG. 4: Correlation length ξ_6 (a) and Frank constant F_A (b) as a function of the inverse temperature. ξ_6 diverges at Γ_i and F_A at Γ_m . In between the system shows hexatic symmetry. The solid lines are fits to eq. (3) and (5) resulting in critical exponents $\nu = 0.5 \pm 0.03$ and $\bar{\nu} = 0.35 \pm 0.02$ respectively. Triangles are shifted by 0.1Γ for clarity.

be the transition temperature T_i . For $\Gamma < \Gamma_i$, F_A should jump to zero which is not completely reproduced. We note that since η_6 is not well defined in the isotropic fluid, it becomes problematic to extract F_A from eqn. (4) below Γ_i . At Γ_m , at the hexatic $\rightarrow$ crystalline transition, F_A must diverge which indeed it does. This divergence can be identified with the divergence of the square of the translational correlation length ξ_+ [4]

$$F_A(\Gamma)/k_B T \sim \xi_+^2 \sim \exp\left(\frac{2c}{|1/\Gamma - 1/\Gamma_m|^{\bar{\nu}}}\right), \quad (5)$$

where c is again a constant and $\bar{\nu} = 0.36963$. Fitting the values of F_A to the expression in eqn. (5) in the range $57.5 < \Gamma < 61$ we obtain $\bar{\nu} = 0.35 \pm 0.02$ and $\Gamma_m = 61.3 \pm 0.4$ as an upper threshold. Again triangles represent evaluation of our data in sub-windows of variable size (same sizes as above). The finite size effect for $\Gamma = 57.0$ is negligible. Close to Γ_m it increases but the values saturates for $\Gamma = 59.1$ and $\Gamma = 60.8$ and remain within the error-bars for the biggest sub-windows.

In conclusion, we have checked quantitatively the change of quasi-long-range to short-range orientational order and extracted the correlation length ξ_6 in the isotropic fluid and Frank's constant F_A in the hexatic phase from trajectories of a 2d colloidal system. We find a hexatic $\leftrightarrow$ isotropic liquid transition at $\Gamma_i = 57.5 \pm 0.5$. Three observations support this result: (i) the change of the distance dependence of $G_6(r)$ (Fig. (3)), (ii) the condition $F_A(\Gamma_i) = 72/\pi$ for Frank's constant and (iii) the divergence of ξ_6 . For the transition hexatic $\leftrightarrow$ crystal F_A diverges at Γ_m . Both divergencies (extracted from just one correlation function) lead to critical exponents that are in good agreement with the KTHNY-theory. The measurements for melting and freezing support each other; so we may conclude that there is no hysteresis effect of the phase-transitions. At the two transitions, the order parameters are observed to change continuously (within the resolution of $\Gamma \propto 1/T$); no indication of a phase-separation (as for example strong fluctuations of the order parameters) has been found [30] as has been reported by [17, 21]. So we believe that in our system - having a well-defined, purely repulsive pair-potential and a confinement to 2D that is free of any surface roughness - the transitions are second order.

In [31, 32] we verified that the Young's modulus becomes 16π at T_m . We have now checked that F_A takes the value $72/\pi$ at T_i . These two findings together confirm the two-stage KTHNY melting scenario with its underlying microscopic picture of breaking the translational symmetry by dislocation-pair- and orientational symmetry by disclination-pair-unbinding.

P. Keim gratefully acknowledges the financial support of the Deutsche Forschungsgemeinschaft.

-
- [1] J. Kosterlitz and D. Thouless, J. Phys. C, **6**, 1181 (1973).
 - [2] A.P. Young, Phys. Rev. B, **19**, 1855 (1979).
 - [3] B.I. Halperin and D.R. Nelson, Phys. Rev. Lett. **41**, 121 (1978).
 - [4] D.R. Nelson and B.I. Halperin, Phys. Rev. B **19**, 2457 (1979).
 - [5] S.T. Chui, Phys. Rev. B **28**, 178 (1983).
 - [6] H. Kleinert, Phys. Lett. **95A**, 381n (1983).
 - [7] M.A. Glaser and N.A. Clark, Adv. Chem. Phys. **83**, 543 (1993).
 - [8] Y. Lansac, M.A. Glaser, and N.A. Clark, Phys. Rev. E **73**, 041501 (2006).
 - [9] K. Chen, T. Kaplan, and M. Mostoller, Phys. Rev. Lett. **74**, 4019 (1995).
 - [10] F.L. Somer, G.S. Canright, T. Kaplan, K. Chen, and M. Mostoller, Phys. Rev. Lett. **79**, 3431 (1997).
 - [11] C.H. Mak, Phys. Rev. E, **73**, 065104(R) (2006).
 - [12] A. Jaster, Phys. Rev. E **59**, 2594 (1999).
 - [13] S.Z. Lin, B. Zheng, and S. Trimper, Phys. Rev. E. **73**, 066106 (2006).
 - [14] C.A. Murray and D.H. Van Winkle, Phys. Rev. Lett, **58**, 1200 (1987).
 - [15] Y. Tang, A.J. Armstrong, R.C. Mockler, and W.J. O'Sullivan, Phys. Rev. Lett, **62**, 2401 (1989).
 - [16] R.E. Kusner, J.A. Mann, J. Kerins, and A.J. Dahm, Phys. Rev. Lett, **73**, 3113 (1994).
 - [17] A.H. Marcus and S.A. Rice, Phys. Rev. Lett, **77**, 2577 (1996).
 - [18] K. Zahn and G. Maret, Phys. Rev. Lett, **85**, 3656 (2000).
 - [19] A.V. Petukhov, D. van der Beek, R.P.A. Dullens, I.P. Dolbnya, G.J. Vroege, and H.N.W. Lekkerkerker, Phys. Rev. Lett, **95**, 077801 (2005).
 - [20] R.A. Segalman, A. Hexemer, R.C. Hayward, and E.J. Kramer, Macromolecules, **36**, 3272 (2003).
 - [21] D.E. Angelescu, C.K. Harrison, M.L. Trawick, R.A. Register, P.M. Chaikin, Phys. Rev. Lett. **95**, 025702 (2005).
 - [22] R. Seshadri and R.M. Westervelt, Phys. Rev. B. **46**, 5142 (1992).
 - [23] P.M. Reis, R.A. Ingale, and M.D. Shattuck, Phys. Rev. Lett. **96**, 258001 (2006).
 - [24] X.H. Zheng and R. Grieve, Phys. Rev. B. **73**, 064205 (2006).
 - [25] P. Dimon, P.M. Horn, M. Sutton, R.J. Birgeneau, and D.E. Moncton, Phys. Rev. B, **31**, 437 (1985).
 - [26] D. Li and S.A. Rice, Phys. Rev. E. **72**, 041506 (2005).
 - [27] P. Keim, G. Maret, U. Herz, and H.H. von Grünberg, Phys. Rev. Lett. **92**, 215504 (2004).
 - [28] The segments will merge to rings, if the systems size tends towards infinity.
 - [29] The upper value is motivated by the maximum value of the histogram of distances and $r/a = 0$ is excluded in the algebraic case to avoid the singularity. The histogram over distances is used as statistical weight of the fit-function taking into account the different frequency of occurrence of the data-points in the minima and maxima of $G_6(r)$.
 - [30] We indeed do see a local clustering of dislocations as already noted by [14, 15] in the hexatic phase close to Γ_i (which is not too surprising for dislocations with finite density having an attractive interaction) but this averages out in G_6 if the field of view is big enough.

- [31] H.H. von Grünberg, P. Keim, K. Zahn, and G. Maret, Phys. Rev. Lett. **93**, 255703 (2004).
- [32] J. Zanghellini, P. Keim, and H.H. von Grünberg, J. Phys. Cond. Mat. **17**, 3579 (2005).